

HAMILTON, BROOK, SMITH & REYNOLDS, P.C.

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under
37 C.F.R. 1.53(b))

Attorney Docket No.

1314.1058-001

First Named Inventor or
Application Identifier

Timothy P. Tully

Express Mail Label No.

EL290726222US

Title of
Invention

Gene Chip Technology For Determining Memory Genes

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents
Box Patent Application
Washington, D.C. 20231

1. ☐ Fee Transmittal Form
(Submit an original, and a duplicate for fee processing)
2. ☒ Specification **Total Pages 50**
(preferred arrangement set forth below)
 - Descriptive title of the invention
 - Cross References to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to microfiche Appendix
 - Background of the Invention
 - Summary of the Invention
 - Brief Description of the Drawings
 - Detailed Description
 - Claim(s)
 - Abstract of the Disclosure
3. ☒ Drawing(s) (35 U.S.C. 113) **Total Sheets 9**
☐ Formal ☒ Informal
4. ☐ Oath or Declaration/POA **[Total Pages []]**
 - a. ☐ Newly executed (original or copy)
 - b. ☐ Copy from a prior application (37 C.F.R. 1.63(d))
(for continuation/divisional with Box 17 completed)
[NOTE Box 5 below]
 - i. ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting
inventor(s) named in the prior
application, see 37 C.F.R. 1.63(d)(2)
and 1.33(b).
5. ☐ Incorporation By Reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a
copy of the oath or declaration is supplied under Box 4b, is
considered as being part of the disclosure of the accompanying
application and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program (Appendix)
7. ☐ Nucleotide and/or Amino Acid Sequence Submission
(if applicable, all necessary)
 - a. ☐ Computer Readable Copy
 - b. ☐ Paper Copy (identical to computer copy)
[] Pages
 - c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & documents)
9. ☐ 37 C.F.R. 3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS
Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
14. ☐ Small Entity Statement(s) ☐ Statement filed in prior application,
status still proper and desired
15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)
16. ☐ Other: _____

17. If a **CONTINUING APPLICATION**, check appropriate box and supply the requisite information:

☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.:

Prior application information: Examiner:

Group Art Unit:

18. CORRESPONDENCE ADDRESS

NAME	Richard W. Wagner, Esq.				
	HAMILTON, BROOK, SMITH & REYNOLDS, P.C.				
ADDRESS	Two Militia Drive				
CITY	Lexington	STATE	MA	ZIP CODE	02421-4799
COUNTRY	USA	TELEPHONE	(781) 861-6240	FAX	(781) 861-9540

Signature	<i>Helen Lee</i>	Date	<i>March 10, 2000</i>
Submitted by Typed or Printed Name	Helen Lee	Reg. Number	39,270

-1-

Date: MARCH 10, 2000 Express Mail Label No. EL290726222US

Inventors: Timothy P. Tully, Joshua I. Dubnau, Michael Davis,
Jan Mous and Ulrich Certa
Attorney's Docket No.: 1314.1058-001

GENE CHIP TECHNOLOGY FOR DETERMINING MEMORY GENES

GOVERNMENT SUPPORT

The invention was supported, in whole or in part, by the following grants from
5 the National Institutes of Health: Grant Nos. F32 HD08087-02, P01 HD33098-03,
7R37MH-47840-09 and 5K05MH-00004-26. The United States Government has
certain rights in the invention.

RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application
10 No. 60/124,085, filed March 10, 1999, which teachings are incorporated herein in
entirety by reference.

BACKGROUND OF THE INVENTION

An attribute that many organisms, including humans, possess is memory of past
events. This attribute has been studied for many decades with much information now
15 available that explains many of its ramifications. For example, two basic types of
memory have been identified: transcription-independent memory, which includes short
term memory, and transcription-dependent memory, which includes long term memory.

An heretofore relatively unknown aspect of memory is the identity of genes that contribute to its manifestation. The identity of the genes that contribute to memory formation is just beginning to be explored. Identification of genes associated with memory formation would provide (a) a genetic epidemiology of cognitive dysfunction, (b) diagnostic tools for individuals carrying different allelic forms of these genes (associative with different performance levels for particular forms of cognition) and (c) new targets for drug discovery ultimately to ameliorate various forms of cognitive dysfunction (and particular drugs could be matched to particular forms of cognitive dysfunction by the diagnostic tests). Thus, it would be useful to have techniques available that would identify the genes that are associated with memory formation.

SUMMARY OF THE INVENTION

The present invention is related to Applicants' discovery that the differential effects on memory formation produced by certain experimental protocols can be used to identify genes involved in transcription-dependent memory formation, particularly long term memory formation. The significant difference between any two experimental protocols is in the induction of transcription-dependent memory. The significant difference between any particular two experimental protocols to be compared is the induction of transcription-dependent memory in the experimental group and the absence of transcription-dependent memory in the control group.

Transcription-independent memory includes various "memory phases", such as short-term memory, intermediate-(or middle-) term memory and (in flies) anesthesia-resistant memory. In common to these forms is that pharmacological inhibitors of RNA transcription do not disrupt these memories. Transcription-dependent memory usually is referred to as long-term memory and inhibitors of RNA synthesis block its appearance.

As a result, Applicants' invention relates to methods of identifying a gene or genes involved in transcription-dependent memory (particularly long term memory)

comprising (a) training non-human animals (particularly non-human mammals, other vertebrates and invertebrates) under conditions sufficient to induce transcription-dependent memory formation in the animals; (b) extracting RNA from brain tissue of the animals trained in step (a); (c) synthesizing DNA probes using the RNA extracted in
5 step (b); (d) exposing the DNA probes synthesized in step (c) to microarray chips containing DNA sequences from genes of the genome of the animals under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of the probes to complementary DNA sequences; (e) detecting the signal produced in step (d); and (f)
10 performing a statistical comparison between the signal detected in step (e) and the signal detected in a control.

In one embodiment, the control is obtained according to a method comprising (i) training non-human control animals under appropriate conditions, wherein the conditions are insufficient to induce transcription-dependent memory formation in the
15 control animals; (ii) extracting RNA from brain tissue of the control animals trained in step (i); (iii) synthesizing DNA probes using the RNA extracted in step (ii); and (iv) exposing the DNA probes synthesized in step (iii) to microarray chips containing DNA sequences from genes of the genome of the animals under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray
20 chips, wherein a signal is produced upon hybridization of the probes to complementary DNA sequences. The experimental conditions of step (a) and step (i) constitute an (experimental) treatment pair. The significant difference between the experimental conditions of step (a) and step (i) is in the induction of transcription-dependent memory.

In a second embodiment, the control is obtained according to a method
25 comprising (i) extracting RNA from brain tissue of non-human control animals; (ii) synthesizing DNA probes using the RNA extracted in step (i); and (iii) exposing the DNA probes synthesized in step (ii) to microarray chips containing DNA sequences from genes of the genome of the animals under conditions appropriate for hybridization

of the DNA probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of the probes to complementary DNA sequences. In this embodiment of the control, the control animals are naïve (untrained) animals.

5 As used herein, a control animal is an animal that is of the same species as, and otherwise comparable to (e.g., similar age, sex), the animal that is trained under conditions sufficient to induce transcription-dependent memory formation in that animal.

10 In a particular embodiment, RNA is extracted from the amygdala of trained or control animals. In another embodiment, RNA is extracted from the hippocampus of trained or control animals. In still another embodiment, the signal from hybridized probes is amplified prior to detection. In another embodiment, a statistical comparison is made (performed, conducted) between the signal detected in step (e) and the signal detected in a control that is obtained by training control animals under conditions
15 sufficient to induce transcription-independent memory but not transcription-dependent long term memory.

20 Transcription-dependent memory can be induced using specific experimental conditions. In one embodiment, transcription-dependent memory is induced in a non-human animal using a spaced training protocol for the fear-potentiated startle response. In a second embodiment, transcription-dependent memory is induced in a non-human animal using a shuttle-box avoidance protocol. In a third embodiment, transcription-dependent memory is induced in a non-human animal using a contextual fear conditioning protocol.

25 The invention also relates to a method of identifying a gene or genes involved in transcription-dependent memory in *Drosophila* comprising (a) training *Drosophila* under conditions appropriate to induce transcription-dependent memory formation in the *Drosophila*; (b) extracting RNA from head tissue of *Drosophila* trained in step (a); (c) synthesizing DNA probes using the RNA extracted in step (b); (d) exposing the

DNA probes synthesized in step (c) to microarray chips containing DNA sequences from genes of the *Drosophila* genome under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of the probes to complementary DNA sequences;

5 (e) detecting the signal produced in step (d); and (f) performing a statistical comparison between the signal detected in step (e) and the signal detected in a control.

In a particular embodiment, the control is obtained according to a method comprising (i) training control *Drosophila* under appropriate conditions, wherein the conditions are insufficient to induce transcription-dependent memory formation in the control *Drosophila*; (ii) extracting RNA from head tissue of the control *Drosophila*

10 trained in step (i); (iii) synthesizing DNA probes using the RNA extracted in step (ii); and (iv) exposing the DNA probes synthesized in step (iii) to microarray chips containing DNA sequences from genes of the *Drosophila* genome under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on

15 the microarray chips, wherein a signal is produced upon hybridization of the probes to complementary DNA sequences. The experimental conditions of step (a) and step (i) constitute an (experimental) treatment pair. The significant difference between the experimental conditions of step (a) and step (i) is in the induction of transcription-dependent memory.

20 In a second embodiment, the control is obtained according to a method comprising (i) extracting RNA from head tissue of control *Drosophila*; (ii) synthesizing DNA probes using the RNA extracted in step (i); and (iii) exposing the DNA probes synthesized in step (ii) to microarray chips containing DNA sequences from genes of the *Drosophila* genome under conditions appropriate for hybridization of the DNA

25 probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of the probes to complementary DNA sequences. In this embodiment of the control, control *Drosophila* are naïve (untrained) flies.

As used herein, a control *Drosophila* is a *Drosophila* that is of the same species as, and otherwise comparable to, the *Drosophila* that is trained under conditions sufficient to induce transcription-dependent memory in that *Drosophila*.

In one embodiment of the method of identifying a gene or genes involved in transcription-dependent memory in *Drosophila*, the DNA probes are labeled with a fluorescent marker and the signal is detected using a fluorescence assay. In a particular embodiment, the signal from hybridized probes is amplified prior to detection. In another embodiment, a statistical comparison is performed between the signal detected in step (e) and the signal detected in a control that is obtained by training control *Drosophila* under conditions sufficient to induce transcription-independent memory but not transcription-dependent memory.

Transcription-dependent memory can be induced in *Drosophila* using a spaced training protocol (e.g., spaced training of olfactory Pavlovian conditioning). Transcription-independent memory can be induced in *Drosophila* using a massed training protocol (massed training of olfactory Pavlovian conditioning).

A statistically significant difference in transcript level for a specific gene between animals trained under conditions sufficient to induce transcription-dependent memory and control animals trained under appropriate conditions that are not sufficient to induce transcription-dependent memory identifies that gene as a candidate memory gene (CMG). In a particular embodiment, a statistically significant difference in transcript level between spaced- and massed-trained groups for a specific gene identifies that gene as a candidate memory gene.

A statistically significant difference in transcript level for a specific gene between animals trained under conditions sufficient to induce transcription-dependent memory and naïve (untrained) control animals identifies that gene as a candidate plasticity gene (CPG). In a particular embodiment, a statistically significant difference in transcript level between spaced-trained and untrained groups for a specific gene identifies that gene as a candidate plasticity gene.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic diagram of a training protocol described in Tully *et al.*, *Cell*, 79:35-47 (1994)), which produces Pavlovian olfactory learning in flies.

Figure 2 is a graphic representation of results showing memory retention after spaced or massed training in normal (wild type) flies or spaced training in transgenic
5 hs-CREB2-r flies after induced expression of CREB repressor (see Yin *et al.*, *Cell*, 79:49-58 (1994)). Learning and early memory (cycloheximide insensitive) are normal in transgenic flies. The additional (protein synthesis-dependent LTM) memory normally produced by spaced training is blocked in transgenic flies. This comparison
10 reveals that the only difference between spaced and massed training is the appearance of a transcription-dependent memory after the former.

Figure 3 is a schematic diagram showing the average difference between signal detected for a specific DNA oligonucleotide probe perfectly matched (PM) (complementary) to a specific section of a specific gene and signal detected for that
15 probe mismatched (MM) to that section of the gene as a result of the introduction of a nucleotide sequence error (mutation) in that section of the gene. The average difference between PM and MM pairs, and usually for 20 pairs per gene, is determined by Affymetrix design software analysis (Affymetrix, Inc., Santa Clara, CA). The squares represent microsequences on a microarray chip.

Figure 4A is a scatterplot representation (spaced versus massed) of the mean
20 signal (mean transformed normalized difference) from an N = 10 chips, for each cloned *Drosophila* gene, each hybridized with DNA probes made from RNA extracted from the heads of normal *Drosophila* exposed 24 hours earlier to either spaced or massed training. Each square represents a specific *Drosophila* gene, 1542 of which are
25 contained on each chip. The candidate memory genes are identified by the lighter shaded squares. The location of the C/EBP gene in the plot is indicated in the figure.

Figure 4B is a scatterplot representation (spaced versus massed) showing the statistically significant values from Figure 4A. Each square represents a specific

Drosophila gene, 1542 of which are contained on each chip. The location of the C/EBP gene in the plot is indicated in the figure.

Figure 5 is a bar graph showing the results from a quantitative polymerase chain reaction (QPCR) experiment. The results confirm the differential effect of spaced
5 versus massed training on the C/EBP gene.

Figure 6 is a bar graph showing the effect of an intertrial interval (ITI) between fear conditioning training trials in rats on subsequent long-term memory. The results define massed and spaced protocols and show that memory of fear-potentiated startle is better after spaced-training than after massed-training.

10 Figure 7 is a bar graph showing the effect on long term memory of overexpression of CREB activator in the amygdala of rats. The results show that overexpression of CREB activator in the amygdala enhances (increases) memory of fear-potentiated startle in rats after massed training.

Figure 8 depict molecular maps of the *adfl* genomic region.

15 DETAILED DESCRIPTION OF THE INVENTION

To produce a specific "long-term memory," an animal is subjected to a specific training protocol under controlled, experimental conditions. In Pavlovian conditioning procedures, for instance, two specific stimuli are presented in temporal contiguity to produce "associative learning and memory." One of the two stimuli is designated a
20 "conditioned stimulus" (CS) and the other is designated an "unconditioned stimulus" (US). The US usually is a natural reinforcer that elicits a "unconditioned response" (UR) before training in a "reflexive" manner. With CS-US pairing, a "conditioned response" (CR) begins to appear in response to the CS before (or in the absence of) presentation of the US. After a CR to a specific CS-US pairing is "learned", memory
25 formation thereafter begins.

Memory formation of this specific, experimental experience can exist in two general forms: a transcription-independent form and a transcription-dependent form.

The former includes various "memory phases," such as short-term memory, intermediate-(or middle-) term memory and (in flies) anesthesia-resistant memory. In common to these forms is that pharmacological inhibitors of RNA transcription do not disrupt these memories. The latter form usually is referred to as long-term memory and inhibitors of RNA synthesis block its appearance.

In animal models, various experimental treatments, such as gene mutation, pharmacological blockade, anatomical lesion or specific training protocols, can affect one or more of these types of memories. In particular, some experimental treatments yield normal amounts of transcription-independent memory but do not yield transcription-dependent memory. Such observations constitute the basis of informative DNA chip comparisons. In general, a comparison is made between two experimental protocols; one (experimental group) that is sufficient to induce both transcription-independent and transcription-dependent memories and one that yields only transcription-independent memory (control group). Any detectable differences in transcript levels between these two protocols then can be attributed specifically to a transcription-dependent memory of the experimentally induced learning. These transcripts are referred to herein as "Candidate Memory Genes" (CMGs).

Although experimental conditions are controlled to induce a specific type of learning, other experimentally uncontrolled forms of learning also may take place.

Thus, although a control group may not yield transcription-dependent memory of the specific experimental task, it nevertheless may yield a transcription-dependent memory of an uncontrolled learning experience. One type of such experience is the potential "nonassociative" forms of learning that occur in response to only the CS or US (alone), or in response to CS-US presentations that are not paired temporally (which is the key requirement for "associative learning"). Hence, transcription-dependent "nonspecific" memories may exist in control groups, as defined above. This observation gives rise to a broader class of transcripts involved with "nonspecific" learning, which we refer to as Candidate Plasticity Genes (CPGs). DNA chip comparisons between an experimental

group, as defined above, and naïve (untrained) animals will yield CPGs, along with CMGs.

Behavior-genetic studies in *Drosophila* have established a pair of training protocols with differential effects on memory formation after a Pavlovian odor-shock learning paradigm. Ten training sessions "massed" together (i.e., with no rest interval
5 between sessions) yields maximal learning (acquisition) and transcription-independent memories (not protein synthesis-dependent) (early memories, short-term memory). In contrast, ten training sessions "spaced" (i.e., with a 15-minute rest interval between sessions) yields equivalent levels of learning and transcription-independent memories (early memories), as well as maximal levels of transcription-dependent memory
10 (including protein synthesis-dependent long-term memory (LTM)). LTM requires spaced training; even 48 massed training sessions fails to induce LTM (Tully *et al.*, *Cell*, 79:35-47 (1994)). Protein synthesis-dependent LTM induced by spaced training is blocked completely via overexpression of CREB repressor (Figure 1) (Yin *et al.*, *Cell*, 79:49-58 (1994)). The resulting memory curve after spaced training, where protein
15 synthesis- and CREB-dependent LTM is blocked, is similar to that produced by massed training in normal flies. In contrast, overexpression of CREB activator induces LTM with less training (one training session) or with massed training (Figure 2) (Yin *et al.*, *Cell*, 81:107-115 (1995)). Hence, the induction of LTM is both protein synthesis- and CREB-dependent. These results demonstrate that the only functional difference
20 between spaced and massed training protocols is the appearance of transcription-dependent memory after the former.

This observation forms the basis of a differential screen to identify additional "downstream" genes that are transcriptionally regulated during transcription-dependent memory formation. DNA probes were synthesized using RNA extracted from the heads
25 of spaced- or massed-trained flies according to methods generally known in the art (see, e.g., Sambrook *et al.*, Eds., *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor University Press, New York (1989); and Ausubel *et al.*, Eds., *Current Protocols In Molecular Biology*, John Wiley & Sons, New York (1997)). RNA

was extracted from fly heads as described previously (see, e.g., Drain *et al.*, *Neuron*, 6:71-82 (1991), which is incorporated herein by reference). Spaced- and massed-training of flies were conducted as described previously (see, e.g., Tully *et al.*, *Cell*, 79:35-47 (1994); and Tully and Quinn, *J. Comp. Physiol.*, 157:263-277 (1985), which are incorporated herein by reference). Complementary DNA (cDNA) probe was synthesized from the extracted RNA according to methods generally known in the art (see, e.g., Sambrook *et al.*, Eds., *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor University Press, New York (1989); and Ausubel *et al.*, Eds., *Current Protocols In Molecular Biology*, John Wiley & Sons, New York (1997)).

10 The complex cDNA probe mixture then was hybridized onto microarray chips containing DNA sequences (target DNA sequences) of 1542 *Drosophila* genes (Affymetrix, Inc., Santa Clara, CA; see also, e.g., U.S. Patent No. 5,445,934; and Ramsay, *Nature Biotechnology*, 16:40-44 (1998), which are incorporated herein by reference). In a particular embodiment, the DNA probes are labeled with a detectable

15 marker (e.g., fluorescent marker). The signal from hybridized DNA probes was amplified and detected according to methods generally known in the art (see, e.g., Sambrook *et al.*, Eds., *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor University Press, New York (1989); and Ausubel *et al.*, Eds., *Current Protocols In Molecular Biology*, John Wiley & Sons, New York (1997)). In a particular

20 embodiment, hybridization was detected using a fluorescence assay. A statistical comparison (DNA chip comparison) was made (performed) by comparing the signal detected between spaced- and massed- trained groups.

A sample size and signal transformation algorithm has been determined that improves the statistical reliability to detect small differences in transcript levels between

25 spaced- and massed-trained groups. In a preferred embodiment, it has been determined that a sample size of 10 chips per treatment group for each treatment protocol (i.e., for each gene, 10 chips for spaced-trained group and 10 chips for massed-trained group) improves the statistical reliability to detect small differences in transcript levels between

spaced- and massed-trained groups. In a preferred embodiment, statistical comparison (DNA chip comparison) is made using the following signal transformation algorithm:

1. Determine the "average difference" between signal detected for a set of primer pairs for specific gene. The average difference between perfect matched (PM) and mismatched (MM) signals is determined by Affymetrix design software analysis (Affymetrix, Inc., Santa Clara, CA).

2. Box-Cox Transformation:
Any average difference value below 10 is eliminated. The remaining average difference values for each gene on each chip then are normalized by the overall average difference (across all genes) for that entire chip.

Grand Mean = Overall Avg. Diff. for all chips and all genes/chip

Normalization Factor (for Chip X) = $\frac{\text{Overall Avg. Diff. for Chip X}}{\text{Grand Mean}}$

Norm (Avg. Diff.) = (Norm. Factor for Chip X) \times (Gene Y on Chip X)

- Transformed Avg. Diff. = $\{\ln [\text{norm}(\text{Avg. Diff.})]\} \times 2720.75$

3. Determine the mean and standard error. Compare mean for spaced- and massed-trained flies for each Gene Y using standard t-tests ($\alpha = 0.05$). If $p \leq 0.05$, then the mean signal transformation for a given gene in spaced-trained flies is considered to be statistically different from the mean signal transformation for that gene in massed-trained flies.

Alternatively, statistical comparison (DNA chip comparison) can be made using the following signal transformation algorithm:

1. Determine the "average difference" between signal detected for a set of primer pairs for specific gene. The average difference between perfect matched (PM) and mismatched (MM) signals is determined by Affymetrix design software analysis (Affymetrix, Inc., Santa Clara, CA).

- 5 2. Box-Cox Transformation:
 Any negative average difference for a given gene is zeroed. The "average difference zeroed" (Avg. Diff. 0) for each gene on each chip is normalized by the overall "average difference" for that entire chip.
 - a)
$$\text{Avg. Diff. 0} = \begin{cases} \text{Avg. Diff.}, & \text{if Avg. Diff.} \geq 0, \\ 0, & \text{if Avg. Diff.} < 0 \end{cases}$$

 - b) Normalization:

$$\text{Grand Mean} = \text{Overall Avg. Diff. for all chips and all genes/chip}$$

$$\text{Normalization Factor (for Chip X)} = \frac{\text{Overall Avg. Diff. for Chip X}}{\text{Grand Mean}}$$

- 15 Norm (Avg. Diff. 0) = Normalization Factor (for Chip X) \times (Gene Y on Chip X) (for each gene)

- c)
$$\text{Transformed Avg. Diff.} = \{\ln [\text{norm}(\text{Avg. Diff. 0})]\} \times 2720.75136$$

3. Determine the mean and standard error. Compare spaced mean versus massed mean with a t-test comparison. If $p \leq 0.05$, then the mean signal transformation for a given gene in spaced-trained flies is considered to be statistically different from the mean signal transformation for that gene in massed-trained flies.

Other signal transformation algorithms can be used in performing a statistical comparison (DNA chip comparison) of signal detected between spaced- and massed-trained groups.

With these proprietary approaches, 1542 statistical spaced versus massed
5 training comparisons (Student t-tests) have been determined from *Drosophila*.
Candidate memory genes (CMGs) (not including transposable elements or
mitochondrial genes), which are transcriptionally regulated during transcription-
dependent memory formation, were identified from the DNA chip comparisons
(statistical comparisons) between spaced- and massed-trained groups. Transcripts that
10 are differentially regulated represent those involved specifically in "associative" LTM
induced by a Pavlovian odor-shock learning paradigm.

CMGs from *Drosophila* determined from statistical analysis using the signal
transformation algorithm in which an average difference value below 10 for a given
gene is deleted, from the spaced versus massed comparison (24-hour memory), are
15 presented in Table 1. CMGs from *Drosophila* determined from statistical analysis using
the signal transformation algorithm in which a negative average difference value for a
given gene is zeroed, from the spaced versus massed comparison (24-hour memory), are
presented in Table 2.

TABLE 1:	Statistical Candidate Memory Genes, derived from statistical analysis (delete avg. diff. values < 10), from the spaced versus massed comparison (24-hour memory).
hikaru genki type1 product	
inositol 1,4,5-trisphosphate receptor	
mitochondrial cytochrome c oxidase subunits, ATPase6, 7 tRNAs	
mitochondrial cytochrome c oxidase subunits, ATPase6, 7 tRNAs	
mitochondrial cytochrome c oxidase subunits, ATPase6, 7 tRNAs	
C/EBP gene	
disabled	
molybdenum cofactor (cin)	
dif	
syntaxin1A (syx-1A)	
fsh membrane protein, 7.6 kb mRNA	
defective chorion fc177 (dec-1) gene	
mitochondrial DNA with 12 tRNAs and 7 genes.	
proteasome (PROSA-28.1.1.)	
cysteine-string protein 32 (csp32)	
FTZ-F1 mRNA	
Mov34 protein mRNA	
transcription factor TFIID 230 kda subunitnt	
croc/FD1=crocodile	
CS-5 pheromone-binding protein homolog OS-E mRNA	
mago-nashi protein (mgn) gene	
transcription initiation factor TFIID 28 kDa subunit mRNA	
Canton S RNA binding protein La/SS-B (DLA/SS-B) mRNA	
angiotensin converting enzyme precursor (Ance) mRNA	
commissureless (comm) mRNA	
nuclear hormone receptor superfamily member DHR78 (DHR78) mRNA	
larval serum protein 1 beta subunit (Lsp-1b) gene	
cut locus mRNA for homeodomain-containing protein	
DRI class II gene for type I regulatory subunit of cAMP-dependent kinase	
PO gene	

TABLE 1: Statistical Candidate Memory Genes, derived from statistical analysis (delete avg. diff. values < 10), from the spaced versus massed comparison (24-hour memory) (continued).

PP1 13C gene for protein phosphatase 1 13C
 mRNA for 51 kDa protein
 genes mst 355a and 355b for male accessory gland secretory protein
 PP-Y mRNA for protein phosphatase Y (EC 3.1.3.)
 anon-66Da, Minute(3)66D and anon-66Db genes
 mRNA for 5HT-dro2A receptor (serotonin receptor).
 eye color protein (garnet) mRNA
 sparkling protein (spa) mRNA
 kinesin-related protein (costal-2) mRNA
 Fuzzy (fuzzy) mRNA
 chitinase (CHT4) gene
 mRNA for rab11
 neu=neuralized mRNA
 anachronism Genomic/mRNA
 TART-B1 transposon putative single-stranded nucleic acid bind
 sodium channel protein (para) gene, exons 9,10,11,12
 mod2.2 (mod(mdg4)) mRNA
 tyrosine kinase mRNA
 leucine-rich repeat/Ig transmembrane protein KEK1 precursor
 glilectin mRNA
 Deformed epidermal autoregulatory factor-1 (Deaf1) mRNA
 Lozenge (lz) mRNA
 hook (hook) mRNA
 SH2/SH3 adaptor protein (Dock) mRNA
 retinoid- and fatty acid-binding glycoprotein mRNA
 blown fuse protein (blow) mRNA
 GCR 101 mRNA
 mRNA for metabotropic glutamate receptor
 mRNA for ladybird late homeodomain transcription factor
 mRNA for putative mitochondrial protein, partial
 colt gene
 mRNA for nuclear protein SA

TABLE 2: Statistical Candidate Memory Genes from *Drosophila*, derived from statistical analysis (negative avg. diff. value is zeroed), from the spaced versus massed comparison (24-hour memory).

hikaru genki type1 gene
 tyrosine kinase hopscotch gene
 syntaxin-1A (syx-1A) gene
 proteasome (PROSA-28.1.1.) gene
 P-glycoprotein (MdR 49) gene
 Mov34 protein gene
 POU domain protein (pdm-1) gene
Drosophila melanogaster epidermal growth factor-like protein (spitz) gene
 (clone 10B-1) germ cell-less protein (gc11) gene
 mago-nashi protein (mgn) gene
 serotonin transporter gene
 transcription initiation factor TFIID 28 kDa subunit gene
 ribonucleoside-diphosphate reductase large subunit gene
 nudel (ndl) gene
 bithorax complex (BX-C) gene cluster
 commissureless (comm) gene
 DNA polymerase gamma gene
 larval serum protein 1 beta subunit (Lsp-1b) gene
 hdl cuticle gene cluster
 alpha-methyldopa hypersensitive gene 1(2)amd gene
 Shaker ShB gene
 DNA-binding protein Elf1 gene
 chorion protein s16 gene
 Adh and Adh-dup genes
 E2F gene
 51 kDa protein gene
 caupolican homeoprotein gene
 5HT-dro2A receptor gene
 odorant binding protein LUSH (lush) gene
 eye color protein (garnet) gene
 adenylyl cyclase isoform DAC9 gene

TABLE 2: Statistical Candidate Memory Genes from Drosophila, derived from statistical analysis (negative avg. diff. value is zeroed), from the spaced versus massed comparison (24-hour memory) (continued).

kinesin-related protein (costal-2) gene
chitinase (CHT1) gene
canoe gene
rab11 gene
imitation-SWI protein (ISWI) gene
receptor guanylyl cyclase (DGC1) gene
tumor supressor (warts) gene
kinesin-like protein (KLP4) gene
myosin-IA gene
cytoplasmic basic protein (deltex) gene
Ca/calmodulin-dependent nitric oxide synthase (NOS) gene
sodium channel protein (para) gene
CKII alpha subunit interactor 1 (CKIIalpha-I1) gene
leucine-rich repeat/Ig transmembrane protein KEK1 precursor (kek1) gene
geranylgeranyl transferase beta-subunit type I (beta GGT-I) gene
hook (hook) gene
SH2/SH3 adaptor protein (Dock) gene
RNA-binding protein lark (lark) gene
retinoid-and fatty acid-binding glycoprotein gene
Dreg-2 protein gene
transcription factor dMax gene
non-histone chromosomal protein Prod (prod) gene
blown fuse protein (blow) gene
orb gene
angel gene
ladybird late homeodomain transcription factor gene

The statistical procedures described above only suggest "statistical candidates." A fundamental aspect of the statistical methods employed (as well as other such methods) is that "false positive" and "false negative" candidates are obtained along with the "true positives." Hence, an independent method of detecting experience-dependent changes in gene transcription must be applied to the "statistical candidates." Such independent methods include Northern blot analysis, quantitative polymerase chain reaction (QPCR) and RNase protection assays, and can be used to confirm the statistical

candidates identified. The quantitative analyses of these data also are subject to false positive and false negative results.

Minor changes in the statistical methods herein can yield a different set of "statistical candidates". Often times, more than one type of data transformation is
5 sufficient to yield a normalized distribution of difference scores. Each data transformation used, however, will yield a different set of statistical candidates. All methods of signal detection also must resolve "baseline values", which are too low for accurate detection. Setting such values to "zero" is one way to deal with this difficulty. Another way is to eliminate such values from the data set (e.g., eliminate values of less
10 than 10, for example).

Chip data provide confirmatory information, gene-by-gene, as to which transcripts are involved with memory. Chip data also provide exact coordinated transcriptional response to different stimuli across all gene transcripts. In particular, chip data provide information as to the coordinated effect a gene transcript has on
15 memory.

Most genes in *Drosophila* have been shown to have mammalian homologs, and such is the case for most *Drosophila* genes involved in memory formation (Dubnau and Tully, *Ann. Rev. Neurosci.*, 21:407-444 (1998)). With the growing knowledge that mammalian homologs can be functionally substituted in *Drosophila* for its fly homolog,
20 the present discovery directly implicates the corresponding mammalian homologs.

The differential effects on long-lasting memory produced by spaced versus massed training is a phenomenon widely observed in the animal kingdom. In particular, a spaced-massed differential effect on long-lasting memory recently has been established for the conditioned fear-potentiated startle effect in rats (a mammalian
25 model system). In the fear-potentiated startle paradigm, memory is inferred from an increase in startle amplitude in the presence of a conditioned stimulus (CS) that has been previously paired with footshock. Massed training in rats (4-CS-shock pairings with a 10-second intertrial interval) produces essentially no transcription-dependent

memory whereas spaced training (4 pairings with an 8-minute intertrial interval) produces significant transcription-dependent memory (Figure 6) (Josselyn *et al.*, *Society for Neurosci.*, 24: 926, Abstract 365.10 (1998)). Moreover, overexpression of CREB activator, delivered to the amygdala via viral vector technology, enhances memory from
5 massed training in a manner directly analogous to that observed in *Drosophila* (Figure 7) (Josselyn *et al.*, *Society for Neurosci.*, 24: 926, Abstract 365.10 (1998)). These data demonstrate a CREB-dependent spaced-massed differential with which to identify mammalian CMGs.

Hence, these specific training protocols are expected to yield CMGs in animals,
10 such as mammals, similar to the CMGs identified in *Drosophila*. The term "animal", as used herein, includes mammals, as well as other animals, vertebrate and invertebrate (e.g., birds, fish, reptiles, insects (e.g., *Drosophila* species), *Aplysia*). The terms "mammal" and "mammalian", as used herein, refer to any vertebrate animal, including monotremes, marsupials and placental, that suckle their young and either give birth to
15 living young (eutharian or placental mammals) or are egg-laying (metatharian or nonplacental mammals). Examples of mammalian species include humans and other primates (e.g., monkeys, chimpanzees), rodents (e.g., rats, mice, guinea pigs) and ruminants (e.g., cows, pigs, horses).

To identify the CMGs in non-human animals (particularly non-human mammals,
20 other vertebrates and invertebrates), DNA probes are synthesized using RNA extracted from brain tissues of spaced- or massed-trained non-human animals according to methods generally known in the art (see, e.g., Sambrook *et al.*, Eds., *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor University Press, New York (1989); and Ausubel *et al.*, Eds., *Current Protocols In Molecular Biology*, John
25 Wiley & Sons, New York (1997)). These probes can be labeled with a detectable marker. In a particular embodiment, DNA probes are synthesized using RNA extracted from the amygdala of spaced- or massed-trained animals and, if required, labeled with a detectable marker. A variety of detectable markers and labeling methods are known in the art, including fluorescent, chemiluminescent, biotin, radioactive, enzymatically

detected and immunologically detected markers (see, e.g., Sambrook *et al.*, Eds., *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor University Press, New York (1989); and Ausubel *et al.*, Eds., *Current Protocols In Molecular Biology*, John Wiley & Sons, New York (1997)). RNA is extracted from brain tissues, such as the amygdala, according to methods available in the art. Spaced- and massed-training of animals are conducted using methods generally known in the art (see, e.g., Josselyn *et al.*, *Society for Neurosci.*, 24: 926, Abstract 365.10 (1998); Cassella and Davis, *Physiol. Behav.*, 36:377-383 (1986); Guzowski *et al.*, *Proc. Natl. Acad. Sci. USA*, 94:2693-2698 (1997); Lamprecht *et al.*, *J. Neuroscience*, 17(21):6443-6450 (1997); Bourtchuladze *et al.*, *Cell*, 79:59-68 (1994); and Kogan *et al.*, *Curr. Biol.*, 7:1-11 (1996), which are incorporated herein by reference). This complex probe mixture then is hybridized onto microarray chips containing DNA sequences (target DNA sequences) of genes of the genome of the animals (see, e.g., U.S. Patent No. 5,445,934; and Ramsay, *Nature Biotechnology*, 16:40-44 (1998), which are incorporated herein by reference). The signal from hybridized DNA probes is amplified and detected according to methods generally known in the art (see, e.g., Sambrook *et al.*, Eds., *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor University Press, New York (1989); and Ausubel *et al.*, Eds., *Current Protocols In Molecular Biology*, John Wiley & Sons, New York (1997)). For example, hybridization (the signal from the hybridized probe) can be detected using fluorescence assays or mass spectrometry. Methods using optical fibers, diode array detection, chemiluminescence/luminescence, latex bead agglutination, direct electrical charge change detection (CCD) and piezoelectric readout can also be used. The signal transformation algorithm described above is used to calculate gene expression levels between spaced- and massed-trained groups. A statistically significant difference in transcript level between spaced- and massed-trained groups for a specific gene identifies a candidate memory gene. Statistical comparison (DNA chip comparison) between spaced- and massed- trained groups can be made using the signal transformation algorithms described above.

In addition to statistical comparisons between spaced- and massed-trained groups, CMGs can be identified from DNA chip comparisons (statistical comparisons) between animals trained using other pairs of experimental protocols. The experimental group is trained under conditions sufficient to induce transcription-dependent memory and the control group is trained under conditions that are not sufficient to induce transcription-dependent memory. The significant difference between any two experimental protocols is in the induction of transcription-dependent memory. The significant difference between any two experimental protocols to be compared is the induction of transcription-dependent memory in the experimental group and the absence of transcription-dependent memory in the control group.

Pairs of experimental protocols that primarily differ in the induction of transcription-dependent memory are known the art. For example, a pair of experimental protocols that primarily differ in the induction of transcription-dependent memory can consist of a spaced training protocol and a massed training protocol. In this embodiment, training an animal using a spaced training protocol is sufficient to induce transcription-dependent memory in the animal. Training an animal using a massed training protocol is not sufficient to induce transcription-dependent memory. As another example, a pair of experimental protocols that primarily differ in the induction of transcription-dependent memory can consist of training a normal (wildtype) animal using a shuttle-box avoidance protocol (particularly a one-trial shuttle-box avoidance protocol) and training an animal in which the fornix is surgically lesioned using the shuttle-box avoidance protocol (Taubenfeld *et al.*, *Nat. Neurosci.*, 2(4):309-310 (1999)). In this embodiment, transcription-dependent memory is induced in the normal (wildtype) animal. Transcription-dependent memory is not induced in the animal in which the fornix is surgically lesioned. As a further example, a pair of experimental protocols that primarily differ in the induction of transcription-dependent memory can consist of training an animal using a contextual fear conditioning protocol (particularly a one-trial contextual fear conditioning protocol) and training an animal habituated to the training chamber before contextual fear conditioning using the contextual fear

conditioning protocol (Imprey *et al.*, *Nat. Neurosci.*, 1(7):595-601 (1998)). In this embodiment, transcription-dependent memory is induced in the animal that had not been habituated to the training chamber prior to contextual fear conditioning.

Transcription-dependent memory is not induced in the animal habituated to the training chamber before contextual fear conditioning. Other pairs of experimental protocols can readily be identified by those skilled in the art.

DNA probes are synthesized using RNA extracted from the brain tissues, such as the amygdala and hippocampus, of animals trained using pairs of experimental protocols, as described herein, and, if required, labeled with a detectable marker. The DNA probe mixtures then are hybridized onto microarray chips containing DNA sequences (target DNA sequences) of genes of the genome of the animals. A statistical comparison is made by comparing the DNA chip data between the two experimental protocols using the signal transformation algorithms described above.

The statistical procedures herein can be used to detect differences in transcript levels between trained and untrained (naïve) groups. Accordingly, candidate plasticity genes (CPGs) can be identified from DNA chip comparisons (statistical comparisons) between trained versus untrained (naïve) animals. Transcripts that are differentially regulated in this class will include the CMGs, along with any other genes that are transcriptionally responsive in a "nonassociative" manner to the general training conditions (e.g., presentations of odors, electroshock or any other experiential aspects of the training protocol). Some nonspecific transcriptional responses occur simply when an animal is placed in a novel environment or when the animal is exposed to a stimulus alone or unpaired in time. These transcriptional changes may result from general (nonspecific) increases in neuronal activity or reflect other forms of learning/memory formation that are not related to the general training conditions.

CPGs from *Drosophila* determined from statistical analysis using the signal transformation algorithm in which an average difference value below 10 for a given gene is deleted, from the spaced versus naïve comparison (24-hour memory), are

presented in Table 3. CPGs from *Drosophila* determined from statistical analysis using the signal transformation algorithm in which a negative average difference value for a given gene is zeroed, from the spaced versus naïve comparison (24-hour memory), are presented in Table 4.

bioRxiv preprint doi: <https://doi.org/10.1101/131410>; this version posted November 14, 2017. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

TABLE 3: Statistical Candidate Plasticity Genes, derived from statistical analysis (delete avg. diff. values < 10), from the spaced versus naïve comparison (24-hour memory).

ribosomal protein S6 (rps6)
 hu-li tai shao (hts) mRNA
 ribosomal protein S3/AP endonuclease DNA repair protein mRNA
 gurken gene
 atonal protein mRNA
 cytoplasmic dynein intermediate chain (Cdic) gene
 glyceraldehyde-3-phosphate dehydrogenase-1 gene
 glyceraldehyde-3-phosphate dehydrogenase-2 gene
 50 kDa protein F1 gene
 Dmras85D gene
 serine protease (SER1 and SER2) genes
 myosin light chain 2 (MLC2) mRNA
 carboxylesterase 6 and P (Est-6 and Est-P) genes
 annexin IX mRNA
 proteasome (PROSA-28.1.1.) mRNA
 laminin B2 gene
 octopamine receptor mRNA
 A2 component of diphenol oxidase (Dox-A2) gene
 homolog of RAD6 (DHR6) mRNA
 Mov34 protein mRNA
 glu-prolyl tRNA aminoacyl synthetase mRNA
 G protein-coupled receptor kinase (GPRK-1) mRNA
 optomotor-blind mRNA
 profilin (chickadee) mRNA
 transcription factor IIB (TFIIB) mRNA
 glutathione S-transferase-related protein mRNA
 trypsin-alpha, -beta and -epsilon genes
 trypsin-alpha, -beta and -epsilon genes
 catalase gene
 proteasome subunit (l(3)73Ai) gene

TABLE 3: Statistical Candidate Plasticity Genes, derived from statistical analysis (delete avg. diff. values < 10), from the spaced versus naïve comparison (24-hour memory) (continued).

Canton S pheromone-binding protein-related protein PBPRP-2 mRNA
transcription initiation factor TFIID 28 kDa subunit mRNA
glutathione-dependent formaldehyde dehydrogenase gene
cofilin/actin depolymerizing factor homolog mRNA
N-ethylmaleimide-sensitive fusion protein mRNA
ribosomal protein DL11 mRNA
rfc40 protein, Rop protein (Rop), and small GTP binding protein
PROS-Dm25g gene for proteasome
DmTnC 41C mRNA for troponin-C
WM6 mRNA
mRNA for 40S ribosomal protein S12
mRNA for mitochondrial ATPase synthase
genes mst 355a and 355b for male accessory gland secretory protein
Pgk gene for phosphoglycerate kinase
D.melanogaster ribosomal protein 15a (40S subunit).
eye color protein (garnet) mRNA
cysteine proteinase-1 (CP1) gene
calcium-binding protein (SCP1) mRNA
cytochrome P450 (CYP4D2) gene
alpha NAC (oxen) gene, complete cds; and G76C pseudogene
alpha NAC (oxen) gene, complete cds; and G76C pseudogene
kinesin-related protein (costal-2) mRNA
transcriptional co-repressor SIN3A (Sin3A) mRNA
chitinase (CHT2) gene
mRNA for rab-related protein 4.
mRNA for still life type 1
cell adhesion molecule encoding (nrm) gene
phosphoglycero mutase (Pglym78) gene

[illegible][illegible]

TABLE 4: Statistical Candidate Plasticity Genes, derived from statistical analysis (negative avg. diff. value is zeroed), from the spaced versus massed comparison (24-hour memory).

daughterless protein (da)
steroid receptor (FTZ-F1B)
POU domain protein (pdm-1)
bithorax complex (BX-C)
nuclear hormone receptor superfamily(DHR96)
transcriptional co-repressor SIN3A (Sin3A)
mRNA for histone H4
mRNA for ladybird late homeodomain
RAD6 (DHR6)
putative serine protease (easter)
serine protease (SER1 and SER2) genes
proteasome (PROSA-28.1.1.)
Mov34 protein
trypsin-alpha, -beta and -epsilon
trypsin-alpha, -beta and -epsilon genes
proteasome subunit (l(3)73Ai) gene
20S proteasome alpha subunit PSMA5 gene
snake locus mRNA for serine protease
D.melanogaster PROS-Dm25g gene for proteasome
male accessory gland secretory protein (serpin)
serine protease SER4 precursor (Ser4)
cysteine proteinase-1 (CP1) gene
clone 6 serine protease mRNA
Drosophila melanogaster dishevelled mRNA, complete cds.
Dmr85D gene, exon 3
G protein-coupled receptor kinase (GPRK-1) mRNA
rfc40 protein, Rop protein (Rop), and small GTP binding protein
tyrosine kinase, partial sequence
mRNA for rab-related protein 4
mRNA for rab-related protein 3
GDP dissociation inhibitor homologue (dGDI) mRNA
(zeste-white 4) mRNA
phosphoinositide 3-kinase, Dp110
phosphatase 2A catalytic subunit
fasciclin III mRNA
annexin IX mRNA

TABLE 4:	Statistical Candidate Plasticity Genes, derived from statistical analysis (negative avg. diff. value is zeroed), from the spaced versus massed comparison (24-hour memory) (continued).
alpha-methyldopa hypersensitive gene l(2)amd GS2 mRNA for glutamine synthase GS1 mRNA for glutamine synthase mRNA for dopamine receptor eye color protein (garnet) mRNA, clathrin like cell adhesion molecule encoding (nrm) gene Nrv 2.2 neuron surface antigen 2 (Nrv2) mRNA R vacuolar ATPase subunit A (vha68-2) gene mRNA for DNMDAR-I AcTr66B gene for actin-related protein DmTnC 41C mRNA for troponin-C kinesin-related protein (costal-2) mRNA microtubule associated protein (asp) mRNA cytoplasmic dynein intermediate chain (Cdic) gene	

The present invention provides methods of identifying a gene or genes involved in transcription-dependent memory (particularly long term memory) comprising (a) training non-human animals (particularly non-human mammals, other vertebrates and invertebrates) under conditions sufficient to induce transcription-dependent memory

5 (particularly long term memory) in the animals; (b) extracting RNA from brain tissue (such as from amygdala, hippocampus) of the animals trained in step (a); (c) synthesizing DNA probes using the RNA extracted in step (b); (d) exposing the DNA probes synthesized in step (c) to microarray chips containing DNA sequences from genes of the genome of the animals under conditions appropriate for hybridization of the DNA

10 probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of the probes to complementary DNA sequences; (e) detecting the signal produced in step (d); and (f) performing a statistical comparison between the signal detected in step (e) and the signal detected in a control.

In one embodiment, the control is obtained according to a method comprising (i)

15 training non-human control animals (particularly non-human mammals, other

vertebrates and invertebrates) under appropriate conditions, wherein the conditions are insufficient to induce transcription-dependent memory in the control animals; (ii) extracting RNA from brain tissue of the control animals trained in step (i); (iii) synthesizing DNA probes using the RNA extracted in step (ii); and (iv) exposing the

5 DNA probes synthesized in step (iii) to microarray chips containing DNA sequences from genes of the genome of the animals under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of the probes to complementary DNA sequences. The experimental conditions of step (a) and step (i) constitute an

10 (experimental) treatment pair. The significant difference between the experimental conditions of step (a) and step (i) is in the induction of transcription-dependent memory.

In a second embodiment, the control is obtained according to a method comprising (i) extracting RNA from brain tissue of non-human control animals; (ii) synthesizing DNA probes using the RNA extracted in step (i); and (iii) exposing the

15 DNA probes synthesized in step (ii) to microarray chips containing DNA sequences from genes of the genome of the animals under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of the probes to complementary DNA sequences. In this embodiment of the control, the control animals are naïve (untrained)

20 animals.

As used herein, a control animal is an animal that is of the same species as, and otherwise comparable to (e.g., similar age, sex), the animal that is trained under conditions sufficient to induce transcription-dependent memory formation in that animal.

25 Transcription-dependent memory can be induced using specific experimental conditions. In one embodiment, transcription-dependent memory is induced in a non-human animal using a spaced training protocol for the fear-potentiated startle response. In a second embodiment, transcription-dependent memory is induced in a non-human

animal using a shuttle-box avoidance protocol. In a third embodiment, transcription-dependent memory is induced in a non-human animal using a contextual fear conditioning protocol.

The present invention will now be illustrated by the following examples, which
5 are not to be considered limiting in any way.

EXAMPLES

EXAMPLE 1 Isolation of *nalyot*^{P1}.

An X-linked *PlacW* transposon (Bier *et al.*, *Science*, 240(4854):913-916 (1988)) was mobilized to generate 2,182 transposant strains with independent insertions on the
10 second and third chromosomes (cf. Boynton and Tully, *Genetics*, 131:655-672 (1992); Dura *et al.*, *J. Neurogent.*, 9:1-14 (1993)). Three-hour memory after a single training session of Pavlovian olfactory learning was quantified with one performance index (PI) for each of these transposant strains. N = 4 PIs then were generated for those strains that scored 70% or less of a wild-type parental strains [*w*¹¹¹⁸ (CS10)]. At this stage of
15 the screen, 93 mutant strains yielded mean three-hour memory scores 70% wild-type or less. Each of these candidate mutant strains then was outcrossed for five generations to the parental strain to equilibrate their (heterogeneous) genetic backgrounds. When three-hour memory again was quantified (N = 4 PIs) in these outcrossed strains, only eight of the 93 candidate mutants still yielded mean scores < 70% wildtype. Finally,
20 "task relevant" sensorimotor tasks were assayed in these eight mutant strains. All eight showed normal shock reactivity; four, G_B335 and E_J51, E_J220 and E_S152, showed significantly reduced olfactory acuity (Boynton and Tully, *Genetics*, 131:655-672 (1992); Dura *et al.*, *J. Neurogent.*, 9:1-14 (1993). [G_B335, now named *dare*, has been studied further and shows preferential expression in antenna (Freeman *et al.*,
25 *Development*, 126:4591-4602 (1999)]. The remaining four mutant strains displayed normal sensorimotor responses and were named *latheo* (Boynton and Tully, *Genetics*,

131:655-672 (1992); Pinto *et al.*, *Neuron*, 23:45-54 (1999); Rohrbough *et al.*, *Neuron*, 23:55-70 (1999), *linotte* (Dura *et al.*, *J. Neurogent.*, 9:1-14 (1993); Bolwig *et al.*, *Neuron*, 15:829-842 (1995); Simon *et al.*, *Mech. Dev.*, 76:42-55 (1998), *golovan and nalyot*.

5 EXAMPLE 2 Cloning and Characterization of *nalyot* Genomic Region.

The *PlacW* transposon includes a unique *SacII* restriction site followed by the bacterial origin of replication and ampicillin resistance gene (Figure 8). Digestion of *nalyot* (*nal*) genomic DNA with *SacII* ligation under dilute conditions and bacterial transformation allowed plasmid rescue of a 9.4 kb *SacII* restriction fragment along with
10 flanking DNA from the genomic region. Chromosome in situs and southern blotting experiments verified that this fragment co-mapped to the P-insertion site. The radiolabeled rescue fragments were used to screen one million plaques of a lambda-DashII *Drosophila* Can-S genomic library (Stratagene). Isolation, subcloning and restriction analysis of 10 independent genomic clones led to the construction of a 35 kb
15 map spanning the genomic region around the P-element insertion site (Figure 8).

Intron/exon maps of the *adfl* and *cn20* transcription units are shown in Figure 8. The *nal^{P1}* element (arrow) is inserted within an intron of the *adfl* transcription unit, 147 bp downstream of the splice donor site. The *adfl* gene encodes a transcription factor distantly related to the myb family (England *et al.*, *Proc. Natl. Acad. Sci. USA*,
20 89:683-687 (1992)) and is alternatively spliced into (at least) two mRNAs. i1 and i2 correspond to two potential translation start sites. Two additional introns of about 3.5 kb and 59 bp appear to be spliced constitutively and separate the remainder of the *adfl* transcription unit into 274 bp and 1,013 bp exons. The *cn20* gene is novel and produces a single, unspliced transcript that can encode a 395 amino acid protein. The
25 extent of the genomic deletion in *nal^{le60}* is indicated. Restriction sites: B, BamHI; E, EcoRI; H, HindIII; S, *SacII*.

EXAMPLE 3 Northern Blot Analysis of *adfl* and *cn20* In Mutant *nal^{Pl}* and Wildtype Flies and cDNA Isolation.

Total RNA from whole adult flies, adult heads or adult bodies was isolated with the TriZOL reagent (BRL). The poly(A) fraction was subsequently purified with
 5 oligo(dT) cellulose (Collaborative Research) or magnetized oligo(dT) beads (Dynal). Purified poly(A) RNA was fractionated by formaldehyde-agarose gel electrophoresis and transferred to a ZetaProbe nylon membrane (BioRad) in 10xSSC. The RNA on the dried membrane was fixed by UV-crosslinking at 2,500 ujoules (Stratalinker). For
 10 initial identification of transcript classes, membrane strips were probed overnight with radiolabeled genomic DNA fragments in high stringency Church and Gilbert Buffer, washed extensively and exposed to Kodak BioMax film.

Selected probes were hybridized to two *Drosophila* adult head cDNA libraries, a lambda gt11 bacteriophage adult head library (Salvaterra) and a pJG4-5 plasmid library (Roshbash). From these two libraries, a total of eleven clones, corresponding to two
 15 independent transcription units, were isolated and evaluated by restriction analysis. Ten clones corresponded to the *adfl* transcription unit. Restriction-mapping and sequence analysis of a subset of these revealed a common 3' end processing site and heterogeneity at the 5' end. The 5' heterogeneity reflected the partial splicing of intron 1 (114 bp) and, perhaps, incomplete first strand synthesis. One clone, *cn20*, corresponded to an
 20 independent, neighboring transcription unit.

To quantify the effect of the P-insertion on *adfl* and *cn20* RNA levels, Northern blots derived from *nal^{Pl}* and wildtype heads or bodies were analyzed as above with radiolabeled probes corresponding to *adfl*, *cn20* and a control RNA *rp49*.

Relative to control levels of *rp49* RNA, *cn20* mRNA expression levels in both
 25 heads and bodies were similar in wildtype and mutant flies. In contrast, *adfl* mRNA expression levels were reduced by at least two-fold in mutant heads and bodies.

EXAMPLE 4 Antibody Production.

The entire *ADFl* open reading frame was inserted into the pET30(a) expression vector (Novagen) as a C-terminal fusion. Robust IPTG induction of ADF1 fusion protein was obtained in transformed BL21 bacteria. The majority of induced protein was in the insoluble, inclusion body fraction and this fraction (isolated 3 hours after induction) was enriched nearly 85% for the ADF1 fusion protein. This fraction was washed extensively with PBS and used directly as antigen. Mouse polyclonal and monoclonal antibodies were obtained by standard procedures (Harlow and Lane, *Antibodies. A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Plainview, NY (1998)). Three mice were inoculated with 50 µg of the inclusion body fraction (in complete Freund's adjuvant), then boosted every two weeks with 50 µg of inclusion body fraction in incomplete Freund's adjuvant. All three mice showed robust immune responses; one was sacrificed for hybridoma fusions. Of 800 candidate hybridoma lines, 17 showed a response in ADF1 dot-blot analyses. Subsequent evaluation of the 17 lines by western blotting and immunochemical assays led to the isolation of ten monoclonal lines, including MAb ADF1-8 and MAb ADF1-17.

EXAMPLE 5 Western Blot Analysis of ADF-1 Protein Levels In Mutant *nal^{P1}* and Wildtype Flies.

Frozen heads or bodies were isolated as previously described (Yin *et al.*, *Cell*, 79:49-58 (1994)), and extracts were prepared by homogenizing 100 µl of frozen head powder in 500 µl of RIPA buffer. Protein concentrations were determined by the Bio-Rad protein assay. Protein samples were denatured in standard loading dye, separated by SDS-PAGE and transferred electrophoretically at 100 mA for 2 hours to a nitrocellulose membrane (Bio-Rad). Each membrane was blocked overnight at 4°C in PBST + 5% milk, then incubated for 1-2 hours with primary antibody in PBST + 5% milk. Primary antibodies used were mouse polyclonal sera against *adfl* (1:1000), mouse monoclonal supernatant (MAb Adfl-17) against *adfl* (1:20), mouse monoclonal

supernatant against TBP (1:5), and mouse monoclonal ascites against α -tubulin (1:50,000) (Sigma). The membrane was washed extensively in PBST and incubated for 1-2 hours with HRP-conjugated anti-IgG secondary antibody (Bio-Rad; 1:500). Following extensive washing in PBST, the conjugated products were visualized by enhanced chemiluminescence (Pierce SuperSignal ULTRA Substrate) and autoradiography.

Relative to control levels of two other proteins (TATA-binding protein and α -TUBULIN), ADF1 expression was reduced at least two-fold in mutant flies.

EXAMPLE 6 Comparison of DNA Chip Data Sets Between Wildtype Flies and A Single-Gene Mutant, *nalyot*.

DNA chip data sets between normal (wildtype) flies and a single-gene mutant, *nalyot* were compared. The *nalyot* mutant was shown to have normal memory after massed training, but LTM was not induced by spaced training. Moreover, the *nalyot* mutation was identified as a transposon insertion in the *Adfl* gene, the effect of which is to reduce the amount of *Adfl* transcript and protein in mutant fly heads.

When all baseline values were set to zero and the data then were analyzed, no significant difference between wildtype flies and *nalyot* mutants was detected for the *Adfl* gene. However, when baseline values of 10 or less were eliminated from the analysis (rather than set to zero), a significant effect on *Adfl* transcription was detected, which corroborated the results obtained by Northern analysis, as described in Example 3. This result constitutes a significant verification for the statistical approach in which average difference values below 10 are deleted from the data set.

EXAMPLE 7 Quantitative Polymerase Chain Reaction.

A given group of flies is subjected to a particular training protocol (spaced or massed) and stored in regular food vials after training. Twenty four hours after training, flies from different groups of a given training protocol (spaced or massed) are collected

into a single 50 ml Falcon tube and quick-frozen in liquid nitrogen. The heads of frozen flies are separated from their bodies by vigorous mechanical shaking. Frozen and separated body parts then are sifted through a series of sieves, ultimately to obtain a homogeneous population of fly heads.

- 5 Combined heads from a training group (spaced or massed) are separated into eight groups. Each group of heads then is ground into a powder with mortar and pestal. The powder is transferred to 5 ml of Trizol solution (Gibco) and stored at -70°C overnight.

- 10 The frozen Trizol/ fly powder solution then is thawed. 2 ml of chloroform is added. The mixture is centrifuged at 3,500 rpm for 10 minutes at 4°C . The extracted RNA (in aqueous layer) is decanted to a fresh tube, and 1.4 ml of isopropanol is added.

- For QPCR, aliquots of the above solution are spun at 8,000 rpm for 20 minutes at 4°C . Isopropanol is decanted, and the pellet is washed 1x in 70% ethanol. The pellet is resuspended in 100 μl of H_2O and an equal volume of phenol/chloroform (Gibco) is added. The solution is centrifuged at 14,000 rpm for 5 minutes at 4°C . The top aqueous layer is decanted to a fresh tube. 200 μl of ice-cold ethanol is added, along with 6 μl of 3 M sodium acetate. The solution is incubated at -20°C for at least 20 minutes.

- 20 The solution is then centrifuged at 14,000 rpm for 20 minutes at 4°C . The pellet is resuspended in 20 μl of H_2O and subjected to RQ1 Rnase free DNase (Promega). RNA concentration is determined.

First-strand cDNAs then are synthesized from 1 μg of the DNase-free RNA samples, and the QPCR assay is performed according to Perkin Elmer Biosystems protocols, using the 7700 ABI Prism with "CYBR" cybergreen fluorescence detection.

- 25 **EXAMPLE 8** Statistical Candidate Memory Gene, C/EBP.

The following C/EBP primers, designed by Quantagene (Paris, France), were used in QPCR experiments: 5'-AGACTACCGATGCGAACAAAC-3' (SEQ ID NO:1)

and 3'-GTCCCTGAACTGGTCGTCTA-5' (SEQ ID NO:2), yielding an expected fragment of 221 bp in size.

For C/EBP data, 8 replicated RNA extractions were obtained from approximately 10,000 heads of flies 24 hours after they were subjected to spaced or to massed training. QPCR reactions for each RNA extraction were run in triplicate in accordance with the method outlined in Example 7.

Results for each C/EBP reaction were normalized for RNA amounts against a paired QPCR reaction for TBP β , a control gene which shows no transcriptional changes in these contexts.

Analyzed this way, the mean number of cycles required for C/EBP amplification to reach the critical value was 21.86 ± 0.56 for the spaced group and 23.78 ± 0.56 for the massed group. It took fewer amplification cycles for the spaced group to reach the critical value, indicating a higher concentration of C/EBP at the beginning of the PCR amplification.

These data indicate C/EBP levels were 3.78-fold higher in the spaced training group than in the massed training group (Figure 5). The experimental results confirm that there is a significant difference for C/EBP (called *slow border cells*, *slobo* in *Drosophila*) transcripts isolated from spaced- versus massed-trained flies. Genetic manipulation of C/EBP in mice enhances long-term memory (Sterneck *et al.*, *Proc. Natl. Acad. Sci. USA*, 95(18):10908-10913 (1998)), and molecular manipulation of C/EBP in *Aplysia* blocks long-term facilitation (a cellular substrate for long-term memory of sensitization) (Alberini *et al.*, *Cell*, 76:1099-1114 (1994)). Therefore, confirmation of C/EBP as a CMG herein constitutes a significant verification that this DNA chip approach can be used in identifying transcripts involved in memory.

EXAMPLE 9 Training Apparatus.

Experimentally naïve male Sprague-Dawley rats (300-350 g) were trained and tested in five identical Plexiglas and wire-mesh cages (8 x 15 x 15 cm) housed in a

sound-attenuated chamber (Cassella and Davis, *Physio. Behav.*, 36:377-383 (1986)). The startle-eliciting stimulus was a 105 dB, 50-ms burst of white noise (rise-decay time 5 ms), against a background white noise of 55 dB. A 3.7-s light CS was produced by an 8-W fluorescent lightbulb (rise-decay time, 100 μ s; BCO foot-lamberts intensity). The floor on each cage consisted of four stainless steel bars, through which a 0.5-s 0.6 or 0.3 mA scrambled foot shock could be delivered.

EXAMPLE 10 Behavioral Training Procedures.

On each of two days before all training, rats were placed in the startle chambers and 15 minutes later presented with 15 startle stimuli. On the single training day for Exp. 1, animals were placed in the startle chamber and 5 minutes later received 4 light-shock pairings with one of the following intertrial intervals (ITIs) (3-seconds, 5-seconds, 10-seconds, 15-seconds, 2-minutes or 8-minutes). Five minutes following the last light-shock pairing, animals were returned to their home cages. Paired massed and spaced training were similarly conducted except that the ITI between the 4 light-shock pairings was 10 seconds and 8 minutes, respectively. Explicitly unpaired massed training trials consisted of massed presentations of the light (4, ITI of 10 seconds) followed 4 minutes later by massed presentations of the shock (4, ITI of 10 seconds).

EXAMPLE 11 Long Term Memory Testing.

Forty-eight hours following training, rats were placed in the startle apparatus and received 30 startle-eliciting stimuli alone followed by 60 startle-eliciting stimuli, half of which occurred 3.2 seconds after the onset of the 3.7 seconds light (Light-Noise trial) and half of which were presented in darkness (Noise Alone trial). The order of the two trial types was irregular. All startle stimuli were presented at an interstimulus interval of 30 seconds. Fear-potentiated startle difference scores, used as an index of LTM, were calculated by subtracting the average startle amplitudes produced by the 30 Noise Alone trials from the average startle amplitudes produced by the 30 Light-Noise trials.

EXAMPLE 12 Short Term Memory Testing.

Short term memory (STM) testing was similar to LTM testing except that it occurred 15 or 40 minutes following training. Twenty Noise Alone stimuli were followed by 15 Light-Noise stimuli and 15 Noise Alone stimuli intermixed. Fear-
5 potentiated startle scores were calculated by subtracting the average Noise Alone score from the average Light-Noise score and used as an index of STM.

EXAMPLE 13 Surgery.

Rats were pre-treated with atropine sulfate (0.4 mg/kg, ip), anesthetized with sodium pentobarbital (60 mg/kg, ip) and placed in a standard stereotaxic instrument. A
10 Hamilton microsyringe (10 μ l) mounted in an infusion pump was used for infusions. Bilateral microinjections (2 μ l) were delivered over 10 minutes through a 30 gauge cannulae aimed at the lateral nucleus of the amygdala (coordinates AP = -2.8, L = \pm 5.2, DV = -8.5 below the surface of the skull) or caudate nucleus (co-ordinates AP = +0.2, L = \pm 3.0, DV = -6.0) according to Paxinos and Watson, *The Rat Brain in Stereotaxic*
15 *Coordinates*, Academic, Sydney, Australia (1986). Infusion cannulae were left in place an additional 10 minutes to ensure diffusion.

EXAMPLE 14 Virus Preparation.

CREB and mCREB cDNAs (obtained from M.E. Greenberg, Harvard University) and LacZ were inserted into the HSV amplicon HSV-PrpUC and packed
20 using the helper 5dl 1.2 (Lim *et al.*, *Biotechniques*, 20:460-469 (1996); Keve *et al.*, *Neuroscience*, 79:435-447 (1997)). Virus was purified on a sucrose gradient, pelleted and resuspended in 10% sucrose. The average titer of the recombinant virus stocks was 4.0×10^7 infectious units/ml and was similar for HSV-CREB and HSV-mCREB. Transgene expression was regulated by the constitutive promoter for the HSV
25 immediate-early gene IE 4/5.

EXAMPLE 15 Immunocytochemistry.

Rats were overdosed with chloral hydrate and perfused with 50 ml PBS followed by 250 ml 4% paraformaldehyde in PBS. The brains were cryoprotected and cut on a microtome (40 μ m sections), and immunocytochemistry was performed on free-floating sections. Brains infected with HSV-LacZ were reacted for β -galactosidase and counterstained with neutral red (according to Lim *et al.*, *Biotechniques*, 20:460-469 (1996); Keve *et al.*, *Neuroscience*, 79:435-447 (1997)). Briefly, to detect β -galactosidase activity, brain slices were allowed to react for 2 hours in a solution comprised of 3.1 mM potassium ferrocyanide, 3.1 mM potassium ferricyanide, 20 mM MgCl₂, 0.1 M PBS and 0.2 mg/ml X-gal (Boehringer-Mannheim).

Analysis of transgene expression in brain infected with HSV-CREB was conducted. Briefly, sections were incubated with 1% H₂O₂ and 0.3% Triton-X for 20 minutes, blocked with 1% bovine serum albumin, 2% normal goat serum and 0.3% Triton-X for 30 minutes and incubated with the primary antibody, CREB (1:1000; Upstate Biotechnology, Lake Placid, NY) overnight at 4°C with constant agitation. Sections were incubated with biotinylated goat-anti rabbit IgG secondary antiserum (1:200 dilution; Vector Laboratories, Burlingame, CA) for 2 hours at room temperature. Sections were rinsed and incubated with avidin-biotin peroxidase complex (ABC) reagent (Vector Laboratories). Immunoreactivity was visualized using diaminobenzidine (DAB) reaction.

EXAMPLE 16 Effect of Intertrial interval Between Fear Conditioning Trials On Subsequent Levels of Long Term Memory.

Mean LTM (\pm SEM; assessed as fear-potentiated startle difference scores), assessed 48 hours following training that consisted of 4 light-shock pairings with ITIs of 3 seconds, 5 seconds, 10 seconds, 15 seconds, 2 minutes and 8 minutes (n=10, 10, 10, 5, 5, 15, respectively), varied with different ITIs ($F_{1,49} = 3.04$, $p < 0.05$). The level of LTM is a linear function of the ITI with longer ITIs producing more robust LTM as shown by a significant linear trend ($F_{1,49} = 7.99$, $p < 0.05$). Massed training (3 seconds, 5 seconds,

10 seconds) produces very weak LTM (roughly 50 units). Increasing the rest interval from 10 seconds to 8 minutes yields an increase in LTM as post-hoc Duncan comparisons reveals that 8 min ITI (spaced) produced significantly greater LTM than 10 seconds, 5 seconds and 3 seconds (massed). The results are depicted in Figure 6.

5 EXAMPLE 17 Effect Of Infusion of HSV Vectors Into the Basolateral Amygdala and Extra-amygdala Areas.

Rats infused with HSV-CREB into the basolateral amygdala (n=17) showed significantly greater LTM than rats similarly infused with PBS (n=7), HSV-LacZ (n=10), or HSV-mCREB (n=11) or rats infused with HSV-CREB into brain regions
10 surrounding the basolateral complex of the amygdala (n=8) or into a control region (the caudate; n=5) ($F_{5,52} = 4.99$, $p < 0.001$). Post-hoc analysis revealed that the level of LTM in rats that received HSV-CREB infusion into the basolateral amygdala was significantly higher than all other groups.

Reactivity to footshock was not different for animals given HSV-CREB (n=17),
15 HSV-mCREB (n=11), HSV-LacZ (n=10) or PBS (n=7) infusion into the basolateral amygdala prior to massed training ($F_{3,41} = 1.41$, $p > 0.05$). Mean shock reactivity was assessed by cage displacement for the 200-ms period after each of the 4 footshocks.

Explicitly unpaired conditioning (in which CS (conditioned stimulus) and US (unconditioned stimulus) are not associated) failed to produce LTM for the association
20 in control rats (n=10) and intra-amygdala infusion of PBS (n=5) or HSV-CREB (n=5) did not change this ($F_{2,17} = 0.44$, $p > 0.05$).

Animals that received HSV-CREB 3 days prior to massed training (3d HSV-CREB, n=10) showed greater LTM when re-tested 14 days following infusion than animals similarly treated with HSV-LacZ (3d HSV-LacZ, n=3) or animals given
25 HSV-CREB 14 days prior to massed training and tested 48 h later (14d HSV-CREB, n=4) ($F_{2,14} = 6.05$, $p < 0.05$).

The teachings of all the articles, patents and patent applications cited herein are incorporated by reference in their entirety.

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the invention as defined by the appended claims.

Approved for release

CLAIMS

What is claimed is:

1. A method of identifying a gene or genes involved in transcription-dependent memory comprising the steps of:
 - 5 (a) training non-human animals under conditions sufficient to induce transcription-dependent memory formation in said animals;
 - (b) extracting RNA from brain tissue of said animals trained in step (a);
 - (c) synthesizing DNA probes using the RNA extracted in step (b);
 - 10 (d) exposing the DNA probes synthesized in step (c) to microarray chips containing DNA sequences from genes of the genome of said animals under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of said probes to complementary DNA sequences;
 - 15 (e) detecting the signal produced in step (d); and
 - (f) performing a statistical comparison between the signal detected in step (e) and the signal detected in a control, wherein said control is obtained according to a method comprising the steps of:
 - 20 (i) training non-human control animals under appropriate conditions, wherein said conditions are insufficient to induce transcription-dependent memory formation in said control animals;
 - (ii) extracting RNA from brain tissue of said control animals trained in step (f)(i);
 - (iii) synthesizing DNA probes using the RNA extracted in step (f)(ii);
 - 25 and

- 5 (iv) exposing the DNA probes synthesized in step (f)(iii) to microarray chips containing DNA sequences from genes of the genome of control animals under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of said probes to complementary DNA sequences.
2. The method of Claim 1 wherein said animal is a non-human mammal.
- 10 3. The method of Claim 1 wherein said transcription-dependent memory formation is long term memory formation.
4. The method of Claim 1 wherein transcription-dependent memory formation is induced using a spaced training protocol and the conditions of step (f)(i) are those according to a massed training protocol.
- 15 5. The method of Claim 1 wherein the conditions of step (f)(i) are those sufficient to induce transcription-independent memory formation but not transcription-dependent memory formation.
6. The method of Claim 5 wherein transcription-independent memory formation is induced using a massed training protocol.
- 20 7. The method of Claim 1 wherein transcription-dependent memory formation is induced using a shuttle-box avoidance training protocol, the control animals of step (f)(i) have a surgical lesion of the fornix and the conditions of step (f)(i) are those according to the shuttle-box avoidance training protocol.

8. The method of Claim 7 wherein said non-human animals are non-human mammals.
9. The method of Claim 1 wherein transcription-dependent memory formation is induced using a contextual fear conditioning training protocol, the control
5 animals of step (f)(i) are habituated to the training chamber before training and the conditions of step (f)(i) are those according to the contextual fear conditioning training protocol.
10. The method of Claim 9 wherein said non-human animals are non-human mammals.
- 10 11. A method of identifying a gene or genes involved in transcription-dependent memory comprising the steps of:
- (a) training *Drosophila* under conditions appropriate to induce transcription-dependent memory formation in said *Drosophila*;
 - (b) extracting RNA from head tissue of *Drosophila* trained in step (a);
 - 15 (c) synthesizing DNA probes using the RNA extracted in step (b);
 - (d) exposing the DNA probes synthesized in step (c) to microarray chips containing DNA sequences from genes of the *Drosophila* genome under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a
20 signal is produced upon hybridization of said probes to complementary DNA sequences;
 - (e) detecting the signal produced in step (d); and
 - (f) performing a statistical comparison between the signal detected in step (e) and the signal detected in a control, wherein said control is
25 obtained according to a method comprising the steps of:

- 5
- (i) training control *Drosophila* under appropriate conditions, wherein said conditions are insufficient to induce transcription-dependent memory formation in said control *Drosophila*;
 - (ii) extracting RNA from head tissue of said control *Drosophila* trained in step (f)(i);
 - (iii) synthesizing DNA probes using the RNA extracted in step (f)(ii); and
 - (iv) exposing the DNA probes synthesized in step (f)(iii) to microarray chips containing DNA sequences from genes of the *Drosophila* genome under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of said probes to complementary DNA sequences.
- 10

- 15
12. The method of Claim 11 wherein said transcription-dependent memory formation is long term memory formation.
13. The method of Claim 11 wherein transcription-dependent memory formation is induced using a spaced training protocol and the conditions of step (f)(i) are those according to a massed training protocol.
- 20
14. The method of Claim 11 wherein the conditions of step (f)(i) are those sufficient to induce transcription-independent memory formation but not transcription-dependent memory formation.
15. The method of Claim 14 wherein transcription-independent memory formation is induced using a massed training protocol.

16. A method of identifying a gene or genes involved in transcription-dependent memory comprising the steps of:
- (a) training non-human animals under conditions sufficient to induce transcription-dependent memory formation in said animals;
 - 5 (b) extracting RNA from brain tissue of said animals trained in step (a);
 - (c) synthesizing DNA probes using the RNA extracted in step (b);
 - (d) exposing the DNA probes synthesized in step (c) to microarray chips containing DNA sequences from genes of the genome of said animals under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a
10 signal is produced upon hybridization of said probes to complementary DNA sequences;
 - (e) detecting the signal produced in step (d); and
 - (f) performing a statistical comparison between the signal detected in
15 step (e) and the signal detected in a control, wherein said control is obtained according to a method comprising the steps of:
 - (i) extracting RNA from brain tissue of non-human control animals;
 - (ii) synthesizing DNA probes using the RNA extracted in step (f)(i);
and
 - 20 (iii) exposing the DNA probes synthesized in step (f)(ii) to microarray chips containing DNA sequences from genes of the genome of control animals under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the
microarray chips, wherein a signal is produced upon
25 hybridization of said probes to complementary DNA sequences.
17. The method of Claim 16 wherein said non-human animal is a non-human mammal.

18. The method of Claim 16 wherein said transcription-dependent memory formation is long term memory formation.
19. The method of Claim 16 wherein transcription-dependent memory formation is induced using a spaced training protocol.
- 5 20. The method of Claim 16 wherein transcription-dependent memory formation is induced using a shuttle-box avoidance training protocol.
21. The method of Claim 20 wherein said non-human animal is a non-human mammal.
22. The method of Claim 16 wherein transcription-dependent memory formation is induced using a contextual fear conditioning training protocol.
- 10 23. The method of Claim 22 wherein said non-human animal is a non-human mammal.
24. A method of identifying a gene or genes involved in transcription-dependent memory comprising the steps of:
- 15 (a) training *Drosophila* under conditions appropriate to induce transcription-dependent memory formation in said *Drosophila*;
- (b) extracting RNA from head tissue of *Drosophila* trained in step (a);
- (c) synthesizing DNA probes using the RNA extracted in step (b);
- (d) exposing the DNA probes synthesized in step (c) to microarray chips
- 20 containing DNA sequences from genes of the *Drosophila* genome under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a

signal is produced upon hybridization of said probes to complementary DNA sequences;

(e) detecting the signal produced in step (d); and

(f) performing a statistical comparison between the signal detected in step (e) and the signal detected in a control, wherein said control is

obtained according to a method comprising the steps of:

(i) extracting RNA from head tissue of control *Drosophila*;

(ii) synthesizing DNA probes using the RNA extracted in step (f)(i); and

(iii) exposing the DNA probes synthesized in step (f)(ii) to microarray chips containing DNA sequences from genes of the *Drosophila* genome under conditions appropriate for hybridization of the DNA probes to complementary DNA sequences on the microarray chips, wherein a signal is produced upon hybridization of said probes to complementary DNA sequences.

25. The method of Claim 24 wherein said transcription-dependent memory formation is long term memory formation.

26. The method of Claim 24 wherein transcription-dependent memory formation is induced using a spaced training protocol.

ABSTRACT OF THE DISCLOSURE

5

Pavlovian Learning in Flies




Fig. 1

Calculation of gene-expression level with Affy chi 20 primer pairs per gene

Average difference between PM and MM signal




Fig. 3




Fig. 4A. Scatterplot of mean transformed normalized average differences from an N = 10 chips each hybridized with probes made from RNA extracted from the heads of normal flies exposed 24 hours earlier to either spaced or massed training.

Fig. 4A

3-1058-101

Spaced24

20000

c/ epb

10000

10000

20000

Massed24

Fig. 4B. Scatterplot as above, but now showing only the statistically significant values.

Fig. 4B

QPCR Detection of C/EBP




Fig. 5

Memory of Fear-potentiated Startle Response in Rats
is Better after
Spaced (≥ 8 m) rather than Massed (≤ 10 s) Training




Fig. 6

Overexpression of CREB activator in the Amygdala
Enhances Memory of
Fear-potentiated Startle in Rats after Massed Training




Fig. 7




Fig. 8